(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 8 February 2001 (08.02.2001)

PCT

(10) International Publication Number WO 01/08673 A1

(51) International Patent Classification7: A61K 31/135

(21) International Application Number: PCT/US00/20834

(22) International Filing Date: 31 July 2000 (31.07.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 60/146,778

31 July 1999 (31.07.1999)

(71) Applicants (for all designated States except US): **SMITHKLINE BEECHAM CORPORATION** [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). NPS PHARMACEUTICALS, INC. [US/US]; 420 Chipeta Way, Salt Lake City, UT 84108 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GOWEN, Maxine [GB/US]; 280 S. Roberts Road, Bryn Mawr, PA 19010 (US). SUVA, Larry, J. [AU/US]; 161 Eaton Drive, Wayne, PA 19087 (US). FOX, John [GB/US]; 4594 Stockbridge Lane, Salt Lake City, UT 84117 (US). STROUP, George, B. [US/US]; 303 Brighton Circle, Devon, PA 19333 (US). NEMETH, Edward, F. [US/US]; 4414 Zarahemla Drive, Salt Lake City, UT 84124 (US).

- (74) Agents: SIMON, Soma, G. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).
- (81) Designated States (national): AE, AL, AU, BA, BB, BG, BR, CA, CN, CZ, DZ, EE, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, TZ, UA, US, UZ, VN, YU, ZA.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CALCILYTIC COMPOUNDS

(57) Abstract: Novel methods of treating bone diseases or disorders are provided.

WO 01/08673 PCT/US00/20834

topical formulation contains suitably 0.01 to 5.0% of a compound of Formula (I). The active ingredient may be administered from 1 to 6 times per day, preferably once, sufficient to exhibit the desired activity, as is readily apparent to one skilled in the art.

As used herein, "treatment" of a disease includes, but is not limited to prevention, retardation and prophylaxis of the disease.

5

10

15

20

25

30

Diseases and disorders which might be treated or prevented, based upon the affected cells, include bone and mineral-related diseases or disorders; hypoparathyroidism; those of the central nervous system such as seizures, stroke, head trauma, spinal cord injury, hypoxia-induced nerve cell damage, such as occurs in cardiac arrest or neonatal distress, epilepsy, neurodegenerative diseases such as Alzheimer's disease, Huntington's disease and Parkinson's disease, dementia, muscle tension, depression, anxiety, panic disorder, obsessive-compulsive disorder, post-traumatic stress disorder, schizophrenia, neuroleptic malignant syndrome, and Tourette's syndrome; diseases involving excess water reabsorption by the kidney, such as syndrome of inappropriate ADH secretion (SIADH), cirrhosis, congestive heart failure, and nephrosis; hypertension; preventing and/or decreasing renal toxicity from cationic antibiotics (e.g., aminoglycoside antibiotics); gut motility disorders such as diarrhea and spastic colon; GI ulcer diseases; GI diseases with excessive calcium absorption such as sarcoidosis; autoimmune diseases and organ transplant rejection; squamous cell carcinoma; and pancreatitis.

In a preferred embodiment of the present invention, the present compounds are used to increase serum parathyroid ("PTH") levels in a non-pulsatile manner. Increasing serum PTH levels may be helpful in treating diseases such as hypoparathyroidism, osteosarcoma,, periodontal disease, fracture, osteoarthritis, rheumatoid arthritis, Paget's disease and osteoporosis.

The normal range for intact PTH in humans is about 10 to about 65 pg/ml. Increasing serum PTH may also be useful to prophylactically retard or prevent the onset of a disease. Prophylactic treatment can be performed, for example, on a person with a low serum PTH, or a person without low serum PTH, but where increasing PTH has a beneficial compensating effect. Preferably, the patient has an

Cot of the

by "activity corporate of the activity petrusou/20834

What is claimed is:

- 1. A method of treating a disease or disorder characterized by an abnormal bone or mineral homeostasis which comprises administering to a subject in need of treatment thereof an effective amount of a calcilytic compound in conjunction with
- treatment thereof an effective amount of a calcilytic compound in conjunction with an effective amount of an anti-resorptive agent.
 - 2. A method according to claim 1 wherein the calcilytic compound is selectedfrom the group consisting of:

N-[(2R-Hydroxy-3-[(3-chloro-2-cyano)phenoxy-propyl]-1,1-dimethyl-2-(2-

- 10 naphthyl)ethyl amine hydrochloride;
 - N-[(2R-Hydroxy-3-[(3-chloro-2-cyano)phenoxy-propyl]-1,1-dimethyl-2-(4-methoxyphenyl)ethyl amine hydrochloride;
 - N-[(2R-Hydroxy-3-[(2,3-dichloro)phenoxy-propyl]-1,1-dimethyl-2-(4-methoxyphenyl)ethyl amine hydrochloride;
- N-[(R)-2-Hydroxy-3-[2-cyano-4-[N-methyl-N-[3-carboxyphenyl)sulfonyl]amino]-phenoxy]propyl]-1,1-dimethyl-2-(6-(1,2,3,4-tetrahydronaphthyl)ethylamine; N-[(R)-2-Hydroxy-3-[2-cyano-4-[N-methyl-N-[3-carboxyphenyl)sulfonyl]amino]-phenoxy]propyl]-1,1-dimethyl-2-(Benzothien-3-yl)-ethylamine; N-[(R)-2-Hydroxy-3-[2-cyano-4-[N-methyl-N-[3-carboxyphenyl)sulfonyl]amino]-
- phenoxy]propyl]-1,1-dimethyl-2-(Benzothien-2-yl)-ethylamine;
 N-[(R)-2-Hydroxy-3-[2-cyano-4-[N-methyl-N-[3-carboxyphenyl)sulfonyl]amino]phenoxy]propyl]-1,1-dimethyl-2-(decahydronapthalen-2-yl)ethylamine;
 N-[(R)-2-Hydroxy-3-[2-cyano-4-[N-methyl-N-[3-carboxyphenyl)sulfonyl]amino]phenoxy]propyl]-1,1-dimethyl-4-phenylbutylamine;
- N-[(R)-2-Hydroxy-3-[2-cyano-4-[N-methyl-N-[3-carboxyphenyl)sulfonyl]amino]-phenoxy]propyl]-1,1-dimethyl-4-(2-methoxyphenyl)butylamine;
 N-[2R-Hydroxy-3-[[2-cyano-4-[N-methyl-N-[4-ethylcarboxyphenyl]sulfonyl]-amino]phenoxy]propyl]-1,1-dimethyl-2-(2-napthyl)ethylamine;
 N-[2R-Hydroxy-3-[[2-cyano-4-[N-methyl-N-[3-
- 30 methylcarboxymethoxyphenyl]sulfonyl]amino]phenoxy]propyl]-1,1-dimethyl-2-(2-napthyl)ethylamine;